



Economic
Research
Service

ERS Elsewhere

MAJOR PUBLICATIONS FEATURING OUR RESEARCH

Least-cost Management of Nonpoint Source Pollution: Source Reduction Versus Interception Strategies for Controlling Nitrogen Loss in the Mississippi Basin

Ecological Economics

Volume 37, Issue 2

May 2001

New York, NY: Elsevier
Science

"Least-cost Management of
Nonpoint Source
Pollution"

Pages 183-197

by Marc O. Ribaud, Ralph
Heimlich, Roger Claassen,
and Mark Peters

For more information, contact

Marc O. Ribaud
at mribaud@ers.usda.gov,
Ralph Heimlich at
heimlich@ers.usda.gov, Roger
Claassen at
claassen@ers.usda.gov, or
Mark Peters at
mpeters@ers.usda.gov
<http://www.ers.usda.gov>

Nutrients are a major
source of water pollution
in the United States.

Agriculture is a major
source of nutrients. Two
alternative strategies for
reducing nutrient loads from
cropland are to reduce fertilizer application rates
and to filter nutrients coming off cropland by
restoring wetlands. A simulation analysis of these
two approaches was conducted for the Mississippi
Basin, where nutrients have caused a large zone of
oxygen-deficient, or hypoxic, waters in the Gulf of
Mexico. Fertilizer reductions impose costs on pro-
ducers by potentially reducing yields and forcing an
inefficient mix of inputs, given prices. Costs for
restoring wetlands include land retirement and
restoration costs. Both approaches can increase
costs for consumers by raising crop prices. Because
of the easement and restoration costs of wetlands, a
fertilizer standard is more cost effective than
restoring wetlands for achieving a water quality goal
up to a point. Beyond that point, wetland restora-
tions are more cost-effective.

